

MICRO-MEC

1.8% Glutaraldehyde Monitor Test Strips

Micro-MEC[™] 1.8% Glutaraldehyde Monitor Test Strip is a glutaraldehyde concentration monitor for use with Micro-Cide[™] 28 High Level Disinfectant. Micro-MEC[™] can also be used with other brands of glutaraldehyde solutions. Micro-MEC[™] cannot be used to validate the sterilization or disinfection process.

FEATURES & BENEFITS

- Monitors minimum effect concentrations of 1.8%.
- Ensures Micro-Cide 28 HLD is effective and safe for use.
- Clear visual standards to communicate PASS/FAIL.
- 2 minute read time.
- Compatible for use with other brands of glutaraldehyde solutions.



ORDER INFORMATION ► Item Code: M60054 • Size: 60 Test Strips • Case: 2 Units/Case





1.8% Glutaraldehyde Monitor Test Strips

DIRECTIONS

- 1. If applicable refer to the endoscope reprocessor's user manual for instructions on how to take a sample of the disinfectant use solution.
- 2. Dip the strip into the solution for 2 seconds and remove.
- 3. Remove excess solution by standing test strip upright (on its short edge) on a paper towel. Do not shake or blot the test strip.
- 4. Start timer.
- 5. At exactly 120 seconds, compare the test strip color with the visual standards below.
- 6. If the test strip color corresponds to 'PASS' according to the visual standards, i.e. uniform purple, then the glutaraldehyde level of the solution is above the MEC.
- 7. The presence of any yellow on the pad (apart from the top 2 mm) indicates that the concentration of glutaraldehyde is below the MEC and the solution should be discarded.

WARNING: Do not use Micro-Cide™ 28 HLD Solution beyond the 28 day maximum use life of the solution.

